Technische Universität München Institut für Informatik Prof. Tobias Nipkow, Ph.D. Alexander Krauss, Sascha Böhme

## Logic

## Exercise Sheet 11

## Exercise 11.1

- a) Show that  $\Sigma \subseteq Th(Mod(\Sigma))$  holds for any set of axioms  $\Sigma$ . When does equality hold?
- b) Consider the language consisting of just the single predicate P. Show that  $Cn(\forall x.P(x))$  is complete.
- c) Give an example for a complete theory T and a formula p (which may contain free variables), such that  $T \not\vDash p$  and  $T \not\vDash \neg p$ .
- d) We have shown that a complete theory is always decidable. Give a counterexample for the reverse implication.

Exercise 11.2 Quantifier elimination for dense linear orders

Apply quantifier elimination to the following formulas:

- a)  $\forall x. \ x < a \Rightarrow x < b$
- b)  $\exists x \ y. \ \forall z. \ (z < x \Rightarrow z \le y) \land (z > y \Rightarrow z \ge x)$
- c)  $\exists x. \neg (\forall y. x = y \Rightarrow (y \ge a \lor y \ge b \lor y \le c))$

Exercise 11.3 Dense linear orders with endpoints

Dense linear orders with endpoints  $\infty$  and  $-\infty$  are axiomatizable by:

$\forall x \ y. \ x = y \lor x < y \lor y < x$	$\forall x. y. x < y \Rightarrow \exists z. x < z \land z < y$
$\forall x \ y \ z. \ x < y \land y < z \Rightarrow x < z$	$\forall x. \ x < \infty \lor x = \infty$
$\forall x. \ \neg(x < x)$	$\forall x\infty < x \lor x = -\infty$

Show that this theory admits quantifier elimination by describing an algorithm.

## Exercise 11.4

The Compactness Theorem for first-order logic states that a countable set of formulas has a model iff all its finite subsets have a model.

Using compactness, show that if a theory is finitely axiomatizable, any countable axiomatization of it has a finite subset that axiomatizes the same theory. In other words, if  $Cn(\Gamma) = Cn(\Delta)$  with  $\Delta$  finite, then there is a finite  $\Gamma' \subseteq \Gamma$  with  $Cn(\Gamma') = Cn(\Gamma)$ .